

gr-qc/0506089
 UFIFT-QG-05-04
 CRETE-05-15

A MEASURE OF COSMOLOGICAL ACCELERATION

N. C. Tsamis[†]

*Department of Physics, University of Crete
 GR-710 03 Heraklion, HELLAS.*

R. P. Woodard[‡]

*Department of Physics, University of Florida
 Gainesville, FL 32611, UNITED STATES.*

ABSTRACT

We propose quantifying the quantum gravitational back-reaction on inflation with an invariant measure of the local acceleration rather than the expansion rate. Our observable is suitable for models in which there is no scalar inflaton to provide a preferred velocity field with which to define the expansion. As an example, we use stochastic techniques to evaluate the local acceleration at one loop order for Λ -driven inflation in pure quantum gravity.

PACS numbers: 04.30.Nk, 04.62.+v, 98.80.Cq, 98.80.Hw

[†] e-mail: tsamis@physics.uoc.gr

[‡] e-mail: woodard@phys.ufl.edu

- *Introduction:* Everyone who has taught introductory physics is familiar with the confusion students sometimes experience between velocity and acceleration. When a stone is thrown upwards in a uniform gravitational field its acceleration is a negative constant whereas its velocity decreases linearly from positive to negative. If the same process were viewed from the frame of an inertial observer initially moving upwards faster than the stone, the stone's velocity would be always negative but its acceleration would be unchanged. We shall argue that a similar confusion exists in identifying an invariant measure of the quantum gravitational back-reaction on inflation. What one *wants* to do is quantify the tendency for quantum processes to decelerate the universe, and this is independent of how rapidly the universe may be expanding with respect to some velocity field.

Simple arguments indicate that gravitational back-reaction should slow inflation, either in pure gravity [1] or in scalar-driven inflation [2]. The idea is that inflation rips from the vacuum a continuous stream of virtual long wavelength quanta which are massless and not conformally invariant. The gravitational interactions between these particles induces a negative energy density that gradually increases as more and more of these particles come into causal contact with one another.

Support for these ideas has come from detailed perturbative computations, at 2-loop order in Λ -driven inflation [3] and at 1-loop order in scalar-driven inflation [4]. Unruh quite properly criticized these results on the grounds that they were obtained by regarding the expectation value of the gauge-fixed metric as an observable [5]. One way of resolving his concern is to do the computation in other gauges. That was too difficult for the 2-loop gravitational process but a simple check in a completely different gauge showed no change in the claimed back-reaction for scalar-driven inflation [6].

A much better way of addressing Unruh's criticism is to construct an invariant *operator* that measures the expansion rate even when perturbations are present. One can then compute the expectation value of this operator. A crude scalar measure of the spacetime expansion rate was obtained using the inverse conformal d'Alembertian acting upon unity [7]. This operator can be promoted to an invariant by evaluating it at an physically defined observation point. When the 1-loop expectation value of that invariant was evaluated for a model which had previously seemed to show back-reaction at one loop, the result was zero secular back-reaction [8]. Curiously, the scalar observable contributed nothing to this. When its expectation value was computed in the old, gauge-dependent coordinate systems it continued to show precisely

the old back-reaction effect. The nullification of the effect was entirely due to the corrections needed to define the observation point invariantly.

A major step forward was taken by Geshnizjani and Brandenberger [9], who worked out how to apply the standard measure of expansion [10] to the case of scalar-driven inflation. In this situation the scalar inflaton $\varphi(t, \vec{x})$ provides a preferred 4-velocity field: ¹

$$u^\mu(t, \vec{x}) \equiv -\frac{g^{\mu\nu}(t, \vec{x}) \partial_\nu \varphi(t, \vec{x})}{\sqrt{-g^{\alpha\beta}(t, \vec{x}) \partial_\alpha \varphi(t, \vec{x}) \partial_\beta \varphi(t, \vec{x})}} . \quad (1)$$

In this expression we mean the quantum metric and the full inflaton operator, including its classical background part and the quantum perturbation:

$$\varphi(t, \vec{x}) = \varphi_0(t) + \delta\varphi(t, \vec{x}) . \quad (2)$$

Expression (1) for $u^\mu(t, \vec{x})$ obviously gives a timelike unit vector which is well-defined for small perturbations. By taking one third of its divergence one obtains a scalar measure of the expansion rate. The scalar becomes a full invariant when evaluated at a physically defined observation point. Owing to the spatial homogeneity and isotropy of the state it is only necessary to invariantly fix the time. The preferred coordinate system is one in which the full scalar agrees with its background value. That is, one solves perturbatively for the operator $\tau[\varphi, g](t, \vec{x})$ which enforces the condition:

$$\varphi(\tau(t, \vec{x}), \vec{x}) = \varphi_0(t) . \quad (3)$$

The invariant expansion measure is:

$$\mathcal{H}(t, \vec{x}) \equiv \frac{1}{3} u^\mu{}_{;\mu}(\tau(t, \vec{x}), \vec{x}) = \frac{1}{3} \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} u^\mu) (\tau(t, \vec{x}), \vec{x}) . \quad (4)$$

When the expectation value of (4) was evaluated for models which seemed to show secular back-reaction at 1-loop order, the result was no secular effect [9]. Computations of quantities such as the stress tensor [11] continue to show secular back-reaction at one loop when observables are evaluated at a gauge-fixed coordinate point instead of at the physically defined location

¹Hellenic indices take on spacetime values while Roman indices take on space values. Our conventions are that the metric $g_{\mu\nu}$ has spacelike signature and the curvature tensor equals $R^\alpha{}_{\beta\mu\nu} \equiv \Gamma^\alpha{}_{\nu\beta,\mu} + \Gamma^\alpha{}_{\mu\rho} \Gamma^\rho{}_{\nu\beta} - (\mu \leftrightarrow \nu)$.

(3). It has also been shown that secular back-reaction can seem to occur at 1-loop order in a two-scalar model if one uses the other, spectator, scalar to measure the expansion rate [12].

It seems clear that there is no secular one loop back-reaction in conventional scalar-driven inflation [13].² This actually agrees with the putative physics of the effect. The causative agent is 1-loop particle production so it would have been fortuitous to see a secular gravitational response at the same order. Secular growth can only come as more and more of the produced particles come into interaction with one another and that must be delayed – in perturbation theory – until at least 2-loop order. Scalar models can be contrived which do show an invariantly quantified back-reaction effect at higher loop order [16]. And the problem will not go away because of the enormous potential impact of back-reaction in a realistic model [17, 18, 19, 20].

A particularly interesting model in which to investigate back-reaction is Λ -driven inflation. The Lagrangian is just that of gravity with a positive cosmological constant:

$$\mathcal{L} = \frac{1}{16\pi G} (R - 2\Lambda) \sqrt{-g} \ , \quad (5)$$

and the perturbative background would be de Sitter. One must certainly go to 2-loop order to see back-reaction in this model [21, 22, 23]. On the other hand, the system is uncomplicated by the classical evolution of a scalar inflaton. The classical result for this model is no evolution at all so, if one sees evolution then it is due to quantum gravitational back-reaction; *except* that one still has to invariantly quantify the effect and there is no scalar inflaton to serve as a clock everyone can agree upon. Of course one could simply define an ersatz scalar any number of ways. For example, one could take the invariant volume of the past light-cone, measured from the observation point back to the initial value surface upon which the quantum state is released. This transforms as a scalar, and it certainly increases monotonically in the timelike direction. The trouble is that many such “scalars” can be defined and none of them has the privileged role that the scalar inflaton plays in setting the zero of time for the post-inflationary universe.

- *The de Sitter example:* The ambiguity and the conflicting results it can produce are easy to understand on the classical level in de Sitter spacetime.

²But unconventional models might show it [14, 15].

Suppose we take the “scalar” to be the time t_c of the “closed” co-ordinate system: ³

$$Closed \quad \Longrightarrow \quad ds^2 = -dt_c^2 + \frac{1}{H^2} \cosh^2(Ht_c) d\Omega_3^2, \quad (6)$$

$$-\infty < t_c < \infty, \quad 0 \leq \alpha_i \leq \pi \quad (i = 1, 2), \quad 0 \leq \alpha_3 < 2\pi. \quad (7)$$

It is elementary to compute the velocity field (1) and the expansion rate (4) in this system:

$$Closed \quad \Longrightarrow \quad u_c^\mu = \delta^\mu_0 \quad \Longrightarrow \quad H_c = H \tanh(Ht_c). \quad (8)$$

The integral curves of this velocity field are initially drawing together – so the expansion rate starts negative – but they eventually draw apart, resulting in positive expansion.

Now suppose we adopt as our “scalar” the time t_o of the “open” co-ordinates sub-manifold: ⁴

$$Open \quad \Longrightarrow \quad ds^2 = -dt_o^2 + e^{2Ht_o} \|d\vec{x}\|^2. \quad (9)$$

$$-\infty < t_o < \infty, \quad 0 \leq \|\vec{x}\| < \infty \quad (10)$$

The resulting velocity field (1) and expansion rate (4) are:

$$Open \quad \Longrightarrow \quad u_o^\mu = \delta^\mu_0 \quad \Longrightarrow \quad H_o = H. \quad (11)$$

For the “open” velocity field the expansion rate is a positive constant throughout the sub-manifold. The latter includes some of the very same points for which the expansion rate is *negative* when measured with the velocity field u_c^μ . When this kind of ambiguity exists on the classical background there is little hope of finding a measure of spacetime expansion upon which all observers can agree when quantum perturbations are present.

A little thought leads to the realization that – in attempting to identify a preferred velocity field from which to measure expansion in Λ -driven inflation – we have succumbed to the same confusion as the introductory physics student who seeks a preferred inertial frame. In the latter case, the laws

³De Sitter spacetime has the topology of $S^{D-1} \times \mathfrak{R}$ and it is natural to cover the full manifold by using a co-ordinate system in which the spatial sections are S^{D-1} .

⁴The “open” co-ordinate system covers half of the full de Sitter manifold and has the topology of $\mathfrak{R}^D$.

of physics are phrased in terms of acceleration. In Galilean relativity, the co-ordinates of a given point in two different inertial frames of reference – which are in relative motion with velocity $\vec{V}$ with respect to one another – are related by:

$$\vec{r} = \vec{r}' + \vec{V}t , \quad (12)$$

$$t = t' . \quad (13)$$

The position and velocity at a given point are frame-dependent quantities while the acceleration is not.

So too in quantum gravity, it is really the local cosmological *acceleration* we wish to measure and not the expansion in some velocity field.

• *Acceleration in de Sitter*: In general, the acceleration is obtained from the deviation equation of two infinitesimally close geodesics $\chi^\mu(\tau)$ and $\chi^\mu(\tau) + \delta\chi^\mu(\tau)$:

$$\frac{D^2 \delta\chi^\mu(\tau)}{D\tau^2} = -R^\mu_{\nu\rho\sigma} [\chi(\tau)] \dot{\chi}^\nu(\tau) \delta\chi^\rho(\tau) \dot{\chi}^\sigma(\tau) . \quad (14)$$

The curvature tensor for de Sitter spacetime equals:

$$de\ Sitter \quad \implies \quad R^\alpha_{\beta\gamma\delta} = H^2 \left(\delta^\alpha_\gamma g_{\beta\delta} - \delta^\alpha_\delta g_{\beta\gamma} \right) . \quad (15)$$

We now consider two initially parallel, timelike geodesics with spacelike separation Δ^μ . In synchronous gauge we have:

$$\chi^\mu(\tau) = \tau \delta^\mu_0 , \quad (16)$$

$$\chi^\mu(\tau) + \delta\chi^\mu(\tau) = \tau \delta^\mu_0 + \Delta^\mu(0) , \quad (17)$$

$$g_{\mu\nu} \dot{\chi}^\mu \Delta^\nu = 0 \quad , \quad g_{\mu\nu} \dot{\chi}^\mu \dot{\chi}^\nu = -1 , \quad (18)$$

so that:

$$\frac{d\chi^\mu(\tau)}{d\tau} \equiv \dot{\chi}^\mu = \delta^\mu_0 . \quad (19)$$

For these geodesics, the deviation equation (14) takes the form:

$$\frac{D^2 \Delta^\mu(\tau)}{D\tau^2} = -H^2 \Delta^\mu g_{\rho\sigma} \dot{\chi}^\rho \dot{\chi}^\sigma + H^2 \dot{\chi}^\mu g_{\rho\sigma} \dot{\chi}^\rho \Delta^\sigma , \quad (20)$$

$$= H^2 \Delta^\mu , \quad (21)$$

and a constant positive acceleration is manifest for any point on the manifold: the deviation between initially parallel, timelike and freely falling observers

expands exponentially at all points in de Sitter spacetime. This is a frame invariant statement that characterizes the de Sitter geometry and, consequently, is free of the ambiguities that plague the velocity field.

- *Acceleration in general:* It is convenient to use the freedom under general co-ordinate transformations to bring an arbitrary metric into synchronous gauge:

$$ds^2 = -dt^2 + g_{ij}(t, \vec{x}) dx^i dx^j . \quad (22)$$

The geodesic deviation equation reduces to:

$$\frac{D^2 \Delta^i(\tau)}{D\tau^2} = -R^i{}_{0j0} \Delta^j , \quad (23)$$

$$= -\left(-\frac{1}{2} g^{ik} \ddot{g}_{kj} + \frac{1}{4} g^{ik} g^{lm} \dot{g}_{kl} \dot{g}_{mj} \right) \Delta^j , \quad (24)$$

The cosmological observation we are interested should not depend on the direction of the vector Δ^i , hence we contract into the vector by multiplying with $g_{ij} \Delta^j$. Nor on the magnitude of the vector Δ^i , hence we scale the vector by dividing with its magnitude $g_{rs} \Delta^r \Delta^s$:

$$\frac{g_{ij} \Delta^j}{g_{rs} \Delta^r \Delta^s} \frac{D^2 \Delta^i(\tau)}{D\tau^2} = \frac{1}{g_{rs} \Delta^r \Delta^s} \left(\frac{1}{2} \ddot{g}_{ij} - \frac{1}{4} g^{kl} \dot{g}_{ik} \dot{g}_{jl} \right) \Delta^i \Delta^j , \quad (25)$$

It is the right hand side of the geodesic deviation equation (14) which should provide the cosmological acceleration measurement γ that the observer performs at event x :

$$\gamma(x) \equiv \frac{1}{g_{rs} \Delta^r \Delta^s} \left(\frac{1}{2} \ddot{g}_{ij} - \frac{1}{4} g^{kl} \dot{g}_{ik} \dot{g}_{jl} \right) \Delta^i \Delta^j . \quad (26)$$

- *Acceleration for flat Robertson-Walker spacetimes:* Let us restrict ourselves to the cosmologically interesting homogeneous, isotropic and spatially flat geometries:

$$FRW \implies g_{ij}(t, \vec{x}) \equiv a^2(t) \delta_{ij} . \quad (27)$$

Derivatives of the scale factor $a(t)$ give the velocity (Hubble) parameter $H(t)$ and the deceleration parameter $q(t)$:

$$H(t) \equiv \frac{\dot{a}}{a} , \quad q(t) \equiv -\frac{a\ddot{a}}{\dot{a}^2} = -1 - \frac{\dot{H}}{H^2} . \quad (28)$$

The relevant components of the curvature tensor are:

$$FRW \quad \Longrightarrow \quad R^i_{0j0} = - \left(H^2 + \dot{H} \right) \delta^i_j = q H^2 \delta^i_j , \quad (29)$$

and, therefore, the cosmological acceleration observable γ equals:

$$FRW \quad \Longrightarrow \quad \gamma(t) = -q H^2 = \frac{\ddot{a}}{a} , \quad (30)$$

and it measures the *fractional* local cosmological acceleration.

• *Stochastic Acceleration:* It is simple to evaluate the expectation value of the local acceleration γ at one-loop order for the gravitational action (5) using stochastic techniques [24, 25, 26] under the assumption of no back-reaction. First, we conformally re-scale the metric and express its time derivatives as functions of the re-scaled metric $\tilde{g}_{ij}$:

$$\begin{aligned} g_{ij} \equiv a^2 \tilde{g}_{ij} &\Longrightarrow \dot{g}_{ij} = 2H a^2 \tilde{g}_{ij} + a^2 \dot{\tilde{g}}_{ij} , \\ &\Longrightarrow \ddot{g}_{ij} = \left(2\dot{H} + 4H^2 \right) a^2 \tilde{g}_{ij} + 4H a^2 \dot{\tilde{g}}_{ij} + a^2 \ddot{\tilde{g}}_{ij} . \end{aligned} \quad (31)$$

We also define the Euclidean direction vector:

$$n^i \equiv \frac{\Delta^i}{\sqrt{\Delta^j \Delta^j}} . \quad (32)$$

Our observable (26) becomes:

$$\gamma(x) = H^2 + \dot{H} + \frac{1}{\tilde{g}_{rs} n^r n^s} \left(\frac{1}{2} \ddot{\tilde{g}}_{ij} + H \dot{\tilde{g}}_{ij} - \frac{1}{4} \tilde{g}^{kl} \dot{\tilde{g}}_{ik} \dot{\tilde{g}}_{jl} \right) n^i n^j . \quad (33)$$

Under the assumption of no back-reaction, the re-scaled spatial metric consists of the identity element plus just the transverse traceless field h_{ij}^{TT} of linearized gravitons:

$$No \text{ Back-Reaction} \quad \Longrightarrow \quad \tilde{g}_{ij} = \delta_{ij} + \sqrt{32\pi G} h_{ij}^{TT} , \quad (34)$$

and to linear order in G , the local acceleration operator is:

$$\begin{aligned} \gamma(x) = H^2 + \dot{H} + \left[\frac{1}{2} \sqrt{32\pi G} \ddot{h}_{ij}^{TT} + H \sqrt{32\pi G} \dot{h}_{ij}^{TT} - 8\pi G \dot{h}_{ik}^{TT} \dot{h}_{jk}^{TT} \right. \\ \left. - \left(16\pi G \ddot{h}_{ij}^{TT} + 32\pi G H \dot{h}_{ij}^{TT} \right) h_{kl}^{TT} n^k n^l \right] n^i n^j + O(G^{\frac{3}{2}}) , \end{aligned} \quad (35)$$

Taking the average over all directions and using:

$$n^i n^j \longrightarrow \frac{1}{3} \delta^{ij} \quad , \quad n^i n^j n^k n^l \longrightarrow \frac{1}{15} \left(\delta^{ij} \delta^{kl} + \delta^{ik} \delta^{jl} + \delta^{il} \delta^{jk} \right) \quad . \quad (36)$$

results in the following expression:

$$\begin{aligned} \int d^2 n \, \gamma(x) = & H^2 + \dot{H} - \frac{8\pi G}{3} \dot{h}_{ij}^{TT} \dot{h}_{ij}^{TT} - \frac{32\pi G}{15} \ddot{h}_{ij}^{TT} h_{ij}^{TT} \\ & - \frac{64\pi GH}{15} \dot{h}_{ij}^{TT} h_{ij}^{TT} + O(G^{\frac{3}{2}}) \quad . \end{aligned} \quad (37)$$

Before substituting the stochastic mode expansion for h_{ij}^{TT} in (37) we must drop the terms where two derivatives act on the same field or upon different fields [24, 25, 26]:

$$Stochastic \implies \int d^2 n \, \gamma(x) = H^2 + \dot{H} - \frac{64\pi GH}{15} \dot{h}_{ij}^{TT} h_{ij}^{TT} + O(G^{\frac{3}{2}}) \quad (38)$$

Now from the expansion of the stochastic field in de Sitter spacetime [24, 25, 26]:⁵

$$h_{ij}^{TT}(t, \vec{x}) = \int \frac{d^3 k}{(2\pi)^3} \theta(Ha(t) - k) \sum_{\lambda} \left\{ \frac{H}{\sqrt{2k^3}} e^{i\vec{k} \cdot \vec{x}} \epsilon_{ij}(\vec{k}, \lambda) \alpha_{\vec{k}, \lambda} + c.c. \right\} \quad (39)$$

it is elementary to calculate its two-point function:

$$\begin{aligned} \langle h_{ij}^{TT}(t, \vec{x}) h_{ij}^{TT}(t', \vec{x}) \rangle &= \int \frac{d^3 k}{(2\pi)^3} \theta(Ha(t) - k) \theta(Ha(t') - k) \\ &\quad \times \frac{H^2}{2k^3} \sum_{\lambda} \epsilon_{ij}(\vec{k}, \lambda) \epsilon_{kl}(\vec{k}, \lambda) \quad , \\ &= \frac{2H^2}{16\pi^3} 4\pi \int_H^\infty \frac{dk}{k} \theta(Ha(t) - k) \theta(Ha(t') - k) \quad , \\ &= \frac{H^2}{2\pi^2} \left\{ \theta(t - t') \ln[a(t')] + \theta(t' - t) \ln[a(t)] \right\} \quad , \\ &= \frac{H^3}{2\pi^2} \left[\theta(t - t') t' + \theta(t' - t) t \right] \quad . \end{aligned} \quad (40)$$

The final answer emerges when we use (40) in expression (38):

$$de\ Sitter \implies \left\langle \int d^2 n \, \gamma(x) \right\rangle = H^2 \left[1 - \frac{32}{15\pi} GH^2 + O(G^2) \right] \quad . \quad (41)$$

⁵In (39), *c.c.* denotes the conjugate expression.

Even for GUT-scale inflation the correction is a very small *constant* effect. Because the stochastic formalism correctly captures only the leading infrared logarithms [24, 25, 26], our result (41) is consistent with zero change at one-loop order which is, in turn, consistent with the physics of back-reaction in quantum gravity on de Sitter spacetime.

Acknowledgements

It is a pleasure to acknowledge years of friendly and stimulating discussions on this topic with L. R. Abramo, R. H. Brandenberger, G. Geshnizjani, A. Guth, D. N. Page and W. G. Unruh. This work was partially supported by the European Social fund and National resources YΠEΠΘ-PythagorasII-2103, by European Union grants FP-6-012679 and MRTN-CT-2004-512194, by NSF grant PHY-0244714, and by the Institute for Fundamental Theory at the University of Florida.

References

- [1] N. C. Tsamis and R. P. Woodard, Phys. Lett. **B301** (1993) 351.
- [2] V. F. Mukhanov, L. R. Abramo and R. H. Brandenberger, Phys. Rev. Lett. **78** (1997) 1624, [arXiv:gr-qc/9609026](#).
- [3] N. C. Tsamis and R. P. Woodard, Ann. Phys. **253** (1997) 1, [arXiv:hep-ph/9602316](#).
- [4] L. R. Abramo, R. H. Brandenberger and V. F. Mukhanov, Phys. Rev. **D56** (1997) 3248, [arXiv:gr-qc/9704037](#).
- [5] W. Unruh, “Cosmological long wavelength perturbations,” [arXiv:astro-ph/9802323](#).
- [6] L. R. Abramo and R. P. Woodard, Phys. Rev. **D60** (1999) 044010, [arXiv:astro-ph/9811430](#).
- [7] L. R. Abramo and R. P. Woodard, Phys. Rev. **D65** (2002) 043507, [arXiv:astro-ph/0109271](#).
- [8] L. R. Abramo and R. P. Woodard, Phys. Rev. **D65** (2002) 063515, [arXiv:astro-ph/0109272](#).
- [9] G. Geshnizjani and R. H. Brandenberger, Phys. Rev. **D66** (2002) 123507, [arXiv:gr-qc/0204074](#).
- [10] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-time* (Cambridge University Press, Cambridge, 1975).
- [11] F. Finelli, G. Marozzi, G. P. Vacca and G. Venturi, Phys. Rev. **D69** (2004) 123508, [arXiv:gr-qc/0310086](#).
- [12] G. Geshnizjani and R. H. Brandenberger, JCAP **0504** (2005) 006, [arXiv:hep-ph/0310265](#).
- [13] P. Martineau and R. H. Brandenberger, “The Effects of Gravitational Back-Reaction of Cosmological Perturbations,” [arXiv:astro-ph/0505236](#).
- [14] G. Geshnizjani and N. Afshordi, JCAP **0501** (2005) 011, [arXiv:gr-qc/0405117](#).

- [15] R. H. Brandenberger and J. Martin, Phys. Rev. **D71** (2005) 023504, **arXiv:hep-th/0410223**.
- [16] L. R. Abramo and R. P. Woodard, Phys. Rev. **D65** (2002) 063516, **arXiv:astro-ph/0109273**.
- [17] R. H. Brandenberger, “Back-Reaction of Cosmological Perturbations,” **arXiv:hep-th/0004016**.
- [18] L. H. Ford, “What does Quantum Field Theory in Curved Spacetime Have to Say about the Dark Energy?”, **arXiv:gr-qc/0210096**.
- [19] R. H. Brandenberger, “Back-Reaction of Cosmological Perturbations and the Cosmological Constant Problem,” **arXiv:hep-th/0210165**.
- [20] R. H. Brandenberger and A. Mazumdar, JCAP **0408** (2004) 015, **arXiv:hep-th/0402205**.
- [21] L. H. Ford, Phys. Rev. **D35** (1985) 710.
- [22] F. Finelli, G. Marozzi, G. P. Vacca and G. Venturi, Phys. Rev. **D71** (2005) 023522, **arXiv:gr-qc/0407101**.
- [23] N. C. Tsamis and R. P. Woodard, “Dimensionally Regulated Graviton 1-Point Function in de Sitter,” **arXiv:gr-qc/0506056**.
- [24] A. A. Starobinskiĭ, “Stochastic de Sitter (inflationary) stage in the early universe,” in *Field Theory, Quantum Gravity and Strings*, ed. H. J. de Vega and N. Sanchez (Springer-Verlag, Berlin, 1986) pp. 107-126.
- [25] A. A. Starobinskiĭ and J. Yokoyama, Phys. Rev. **D50** (1994) 6357, **arXiv:astro-ph/9407016**.
- [26] N. C. Tsamis and R. P. Woodard, “Stochastic Quantum Gravitational Inflation,” **arXiv:gr-qc/0505115**.